

COPIKTRA can cause fatal and/or serious toxicities including infections, diarrhea or colitis, cutaneous reactions, and pneumonitis. Contact the healthcare provider on this card **immediately** if your patient reports any signs or symptoms.

+ IMPORTANT SAFETY INFORMATION YOU SHOULD KNOW:

 I am taking COPIKTRA for my treatment of leukemia or lymphoma.

FOR HEALTHCARE PROVIDERS

+ IMPORTANT SAFETY INFORMATION FOR PATIENTS TAKING COPIKTRA™

Patient Name

COPIKTRA Prescriber Name

Prescriber Phone Number


Copiktra™
(duvelisib)

**PATIENT SAFETY
WALLET CARD**



**Show this to any doctor
involved in your care**

+ IMPORTANT SAFETY INFORMATION

FOR THE PATIENT

COPIKTRA can cause these serious side effects that may lead to death:

- Infections
- Diarrhea or inflammation of your intestine (colitis)
- Skin reactions
- Inflammation of the lungs

If you have any of these symptoms, call your healthcare provider **right away**:

- Fever, chills, or other signs of infection
- New or worsening diarrhea
- Stool with mucus or blood
- Severe stomach-area (abdominal) pain
- New or worsening skin rash or other skin reactions, including:
 - Painful sores or ulcers on your skin, lips, or in your mouth
 - Severe rash with blisters or peeling skin
 - Rash with itching
 - Rash with fever
- New or worsening cough or difficulty breathing



IMPORTANT TO REMEMBER: Call your doctor or get emergency medical care right away if you have any of these symptoms! Show this card to any doctor involved in your care!